



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

left there is another, evidently imperfect and apparently partly superimposed upon the remains of a third one belonging to the opposite (under) side of the specimen. If a fourth one was present it is not now apparent.

NEW YORK BOTANICAL GARDEN.

AN ABNORMAL LEAF IN RUMEX

BY S. B. PARISH.




FIG. 3. Abnormal leaf of *Rumex hymenosepalus*, about $\frac{1}{3}$ natural size.

The accompanying figure represents an abnormal leaf of *Rumex hymenosepalus*, having two blades. The superior blade stands more vertically erect than the lower, it is shorter and more crisped, but in other respects the two are alike. Along the midrib the bases of the blades are separated by an interval of about three millimeters. Two adjacent plants were seen, each having fully half of its leaves affected in this manner, but not all to so great an extent as the one figured. On some the secondary blade was present but as a fragment, of greater or less size, or two or three separated fragments. These might occur at any point along the midrib, from its base nearly to its apex, but always of the form and size which the blade would there have

presented, had it been continuous. And in every case, even when most fragmentary, the secondary blades were produced on both sides of the midrib.

It is possible to regard this teratological condition as an instance of foliar peloria. But to this view there are two objections.

The abnormal blades, or fragments of blades, appeared to have originated at the earliest stage of leaf-formation, rather than to have been outgrowths at a later period. Furthermore, a decided disarrangement of the fibrovascular bundles of the midrib suggests a different explanation. Such a condition might result from the cohesion, or rather fusion, of the midribs of two leaves, one superimposed upon the other, the blade of the uppermost being either reduced or fragmentary. But to whatever cause this malformation was due, it is probably one of infrequent occurrence, since none of a like nature is recorded by Masters.

SAN BERNARDINO, CALIFORNIA.

SHORTER NOTES

A NEW MIKANIA FROM CUBA. — *Mikania alba* sp. nov. Glabrous, except the involucre bracts. Primary branches round, striate, secondary ones hexagonal, canaliculate: petioles 4–6 mm. long, slightly winged; leaf-blades coriaceous, lanceolate-ovate, obtuse at the apex, rounded or obscurely cordate at the base, 4–5.5 cm. long, 1.6–1.8 cm. broad, diminishing in the inflorescence to linear or subulate bracts, three-nerved, the nerves deeply impressed in the reticulate-rugose upper side, prominent on the lower, the margin conspicuously revolute: inflorescence bracted, paniculate, ultimately composed of opposite, widely divaricate racemes, with a terminal one; racemes with 8–24 heads; heads opposite below, alternate above, pedicelled, subtended by a subulate bractlet much exceeding the pedicel; involucre bracts oblong-lanceolate, obtuse, 5–6 mm. long, brownish at the center, becoming transparent towards the ciliate margin, as long as or slightly shorter than the corolla; pappus white, equalling the corolla or nearly so, armed with minute spinules; corolla white, tubular-campanulate, scarcely longer than the involucre bracts, with oblong-lanceolate, acute lobes, thrice shorter than the tube, or less; stamens as long as the corolla, rarely exerted, with cylindrical anthers; style-branches widely divaricate, subsequently recurved-coiling, finely tuberculate: achenes glabrous.

Collected in the Sierra Maestra, at an elevation of 3,400 feet, on Jiquarito Mountain, Sevilla Estate, province of Santiago, Cuba, by the writer on September 18, 1906; *no. 516* (type), in herb. New York Botanical Garden.